DEPRESSION SCREENING TEST

Please use the results of this test as a guide and not a diagnosis, only a licensed mental health practitioner can diagnose depression. When completed **total the number of "Yes" responses**.

1.	Have you felt sad or to	earful for a majority of the day for at least the last two weeks? NO
2.	Have you lost interestYES	in activities which you used to enjoy?NO
3.	Is it difficult for you to are tired the next day?YES	of fall asleep most nights to the point that you do not get adequate sleep and NO
4.	Do you feel you sleep least the last two weelYES	a lot more than you should because of feeling fatigued most of the day for at ks? NO
5.	days or feeling the urg	ange in your appetite that has resulted in either not feeling hungry most ge to eat more than usual and has resulted in a change in your weight?NO
6.	•	decreased to the point that normal daily activities seem overwhelming?NO
7.	complicated tasks?	it seems more difficult to stay focused on activities or to concentrate onNO
8.	Have you thought abo	ut suicide or what it would be like if you were not around anymore?NO
9.	Do you find yourself g	etting angry easily or lashing out at people without a valid reason? NO
10		ell you that they are concerned about you because of your feelings of atterns, or your anger?NO
11	. Is it more difficult for easy?	you to make decisions, even regarding simple matters which used to be
	YES	NO

12. Do you cry more	easily than you used to:
YES	NO
13. Do you often cr made?	icize yourself about things you have done in the past or about decisions you have
YES	NO
o .	radness or anger, changes in your sleep pattern, or a lack of motivation and the way of achieving goals or performing work activities?NO
15. Does the future	ook bleak or even hopeless to you?
YES	NO

Response Results:

0 = No depression

1-4 = Mild depression

5-9 = Moderate depression

10-15 = Major depression

For more information, contact **Psychiatric Institute of Washington**

4228 Wisconsin Avenue, NW Washington, DC 20016 (202) 885-5610

www.psychinstitute.com



Disclaimer: This screening is not a diagnostic tool and in no way should take the place of an assessment administered by a mental healthcare professional. If you are feeling unsafe, please contact us or 911 immediately.

CAGE-AID Questionnaire

When thinking about drug use, illegal drug use, and the use of prescription drug use other than prescribed.

Questions

Have you felt that you should cut down on your drinking or drug use?		
	□YES	□NO
Have people	annoyed you by criticizing your drinking o	r drug use?
	□YES	□NO
Have you felt	bad or guilty about your drinking or drug	use?
	□YES	□NO
Have you ove	or had a driple or drugg first thing in the mo	orning to steady your nerves or get rid a hangover?
nave you eve	er had a drink or drugs first thing in the mo	filling to steady your nerves or get rid a nangover?
	□YES	□NO

Scoring: regard one or more "Yes" responses as a positive screen

For more information, contact **Psychiatric Institute of Washington**

4228 Wisconsin Avenue, NW
Washington, DC 20016
(202) 885-5610
www.psychinstitute.com



Disclaimer: This screening is not a diagnostic tool and in no way should take the place of an assessment administered by a mental healthcare professional. If you are feeling unsafe, please contact us or 911 immediately.

Permission for use granted by Richard Brown, M.D.